



Pool or Spa Submersion: Estimated Nonfatal Drowning Injuries and Reported Drownings, 2017 Report



May 2017

Ted Yang
Directorate for Epidemiology
Division of Hazard Analysis
U.S. Consumer Product Safety Commission
4330 East West Highway
Bethesda, MD 20814

~~CPSA (b)(1) CLEARED BY PUBLIC~~
 NO MFPS/PRVTBLRS OR
PRODUCTS IDENTIFIED *RT*
9/22/2017
EXCEPTED BY: PETITION
RULEMAKING ADMIN. PROCG
WITH PORTIONS REMOVED: _____

This analysis was prepared by CPSC staff and has not been reviewed or approved by, and may not necessarily reflect the views of, the Commission.

Executive Summary

- This report presents annual estimates of the number of emergency department-treated, pool- or spa¹-related, nonfatal drownings (submersion injuries²) between 2014 and 2016, and it presents counts of reported pool- or spa-related drownings (submersion fatalities³) between 2012 and 2014 for children younger than 15 years of age. The subset of submersion injuries and fatalities involving children younger than 5 years of age is also provided. Please note that circulation/suction entrapments in pools or spas are presented in a separate document.⁴ It is also important to note that incidents covered by this report were associated with a pool or spa, but the primary cause of the incident was not necessarily the pool or spa product. The period for reported injury and fatality statistics differs due to the lag in fatality reporting. Parents, caregivers, pool operators, state and local officials, and the media are encouraged to visit PoolSafely.gov for vital safety information regarding the prevention of child drownings and drain entrapments in and around pools and spas.

Key findings:

- There were, on average, an estimated 5,900 pool- or spa-related, hospital emergency department (ED)-treated nonfatal drowning injuries each year for 2014 through 2016; and there were 356 pool- or spa-related drownings reported per year for 2012 through 2014, involving children younger than 15 years of age.
- The majority of the estimated ED-treated, nonfatal drowning injuries for 2014 through 2016, and the reported drownings for 2012 through 2014, were associated with pools (versus spas).
- Annual estimates of the number of children who were treated in hospital emergency departments for pool- or spa-related nonfatal drowning injuries in 2016 were not statistically different from those in 2015.
- For children younger than 15 years of age, there were approximately twice as many reported drownings involving male victims than reported drownings involving female victims.
- Seventy-six percent of the reported drownings from 2012 through 2014, and an annual average of 74 percent of the ED-treated nonfatal drowning injuries from 2014 through 2016, involved children younger than 5 years of age.

¹ The term “spa” is used to refer to spas and hot tubs.

² The term “submersion injury” is used in lieu of the term “non-fatal drowning” when comparing or referring to injuries resulting from incidents involving other products or hazards.

³ The term “submersion fatality” is used in lieu of the term “drowning” when comparing or referring to fatalities resulting from incidents involving other products or hazards.

⁴ 2012–2016, “Reported Circulation/Suction Entrapment Incidents Associated with Pools, Spas, and Whirlpool Bathtubs, 2017 Report,” May 2017.

- Children between the ages of 1 and 3 (12 months through 47 months) represented 61 percent of estimated nonfatal drowning injuries for 2014 through 2016, and that age group represented 65 percent of the reported drownings for 2012 through 2014, for victims younger than 15 years.
- For children younger than 15 years old, 43 percent of the victims of estimated ED-treated pool or spa submersion injuries for 2014 through 2016 were admitted to the hospital or treated and transferred to another hospital, compared to 4 percent for ED-treated injuries to children younger than 15 years old involving all consumer products in the CPSC's jurisdiction during the same period.
- Forty-eight percent of the estimated nonfatal drowning injuries for 2014 through 2016, and 77 percent of the reported drownings for 2012 through 2014, involving children younger than 15 years old, occurred at a residence.
- Residential locations dominated reported incidents involving victims younger than 5 years of age (55 percent for nonfatal drowning injuries from 2014 through 2016, and 86 percent for drownings from 2012 through 2014).
- Most reported fatalities from drowning occurred on the day of (68 percent) or within a week of (additional 26 percent) the incident. Only 6 percent of fatal victims younger than 15 survived beyond a week of the incident, and these victims had severe injuries and required intensive medical care before death.
- Approximately 54 percent of reported drownings (annual average of 192) occurred in in-ground pools. Above-ground pools accounted for 19 percent (annual average of 69), with portable pools accounting for 4 percent (annual average of 13) for children younger than 15 years of age.

Emergency Department-Treated Injury Estimates

For 2014 through 2016, an estimated annual average of 5,900 children younger than 15 years of age were treated in U.S. hospital emergency departments (EDs) for nonfatal injuries associated with pool or spa submersions. Estimates are shown in Table 1. Estimates are also provided for injured children younger than 5 years of age and children 5 to 14 years of age.⁵ Injury estimates came from CPSC’s National Electronic Injury Surveillance System (NEISS) data, where sampling weights are used to project the cases from NEISS hospitals to national estimates. The corresponding annual average estimates for the years 2013 through 2015 are 5,600 children younger than 15 and 4,300 children younger than 5 years of age treated in hospital emergency departments for nonfatal drowning injuries related to pools or spas.

Table 1
Estimated Number of Emergency Department-Treated Pool or Spa Nonfatal Drowning Injuries
Children Younger than 5 and 15 Years of Age, 2014-2016

Year	Estimated Emergency Department-Treated Injuries ⁶		
	Younger than 5 Years	5-14 Years	Younger than 15 Years
Average	4,400	1,600	5,900
2016	4,200	1,800	6,000
2015	4,600	1,500	6,000
2014	4,400	1,400	5,800

Source: U. S. Consumer Product Safety Commission: National Electronic Injury Surveillance System. Appendix A details the methodology for data extraction.

The 2016 estimates of children younger than 15 years of age and children younger than 5 years of age who were treated in U.S. hospital emergency departments for pool- or spa-related nonfatal drownings are not statistically different from the 2015 estimates. On average, during 2014 through 2016, 74 percent of children treated in emergency departments for pool- or spa-related, nonfatal drowning injuries were younger than 5 years of age. Children younger than 5 years of age comprised an estimated 76, 76, and 70 percent of the childhood pool- or spa- related treated injuries in 2014, 2015, and 2016, respectively.

⁵ Estimates for children under age 5 and age 5 to 9 may not sum to the under age 15 total due to rounding.

⁶ The estimates are rounded to the nearest hundred.

Table 2 shows the percent of estimates for 2014 through 2016, associated with pool or spa submersions by type of product. Spa-related submersions constitute 3 percent of the estimated number of the pool or spa submersion treated nonfatal drowning injuries for children younger than 15 years of age, and 5 percent of the estimated number of the pool or spa submersion treated nonfatal drowning injuries for children younger than 5 years of age.

Table 2
Percent of Estimated Emergency Department-Treated Pool or Spa Nonfatal Drowning Injuries
Children Younger than 5 and 15 Years of Age by Product Type, 2014-2016

Product Type	Emergency Department-Treated Injury Percentages		
	Younger than 5 Years	5-14 Years	Younger than 15 Years
Pool	95	100	97
Spa	5	0	3

Source: U. S. Consumer Product Safety Commission: National Electronic Injury Surveillance System. Appendix A details the methodology for data extraction.

Table 3 shows the percentage of the estimated number of pool- or spa-related, nonfatal drowning injuries by victim gender. Male children are more frequently treated for pool- or spa-related nonfatal drowning injuries than female children. This is true of all injured children younger than 15 and the subset of children younger than 5 years of age.


Table 3
Percent of Estimated Emergency Department-Treated Pool or Spa Nonfatal Drowning Injuries
Children Younger than 5 and 15 Years of Age by Gender, 2014-2016

Gender	Estimated Emergency Department-Treated Injury Percentages		
	Younger than 5 Years	5-14 Years	Younger than 15 Years
Male	57	67	60
Female	43	33	40

Source: U. S. Consumer Product Safety Commission: National Electronic Injury Surveillance System. Appendix A details the methodology for data extraction.

Figure 1 illustrates the monthly distribution of the percentages of the estimated emergency department-treated, nonfatal drowning injuries for each age group. The months of May, June, July, and August had the largest percentages.


Figure 1
Percent of Estimated Emergency Department-Treated Pool or Spa Nonfatal Drowning Injuries Children Younger than 15 Years of Age by Month of Treatment 2014-2016



Source: U. S. Consumer Product Safety Commission: National Electronic Injury Surveillance System.

Figure 2 plots the percent of the estimated number of ED-treated, nonfatal drowning injuries as a function of the victim's age. Children younger than 1 year of age accounted for 2 percent of the estimated pool- or spa-related, nonfatal drowning injuries. Children between the ages of 1 and 3 years (12 to 47 months) comprised approximately 61 percent of the estimated number of children treated for pool- or spa-related, nonfatal drowning injuries. An additional 11 percent of the estimated childhood pool- or spa-related, nonfatal drowning injuries occurred in children 4 years of age (48 to 59 months). Children ages 5 to 9 and 10 to 14 accounted for 20 and 6 percent, respectively, of the estimated ED-treated pool or spa-related, nonfatal drowning injuries.

Figure 2
Percent of Estimated Emergency Department-Treated Nonfatal Drowning Injuries by Age
Children Younger than 15 Years of Age, 2014-2016



Source: U. S. Consumer Product Safety Commission: National Electronic Injury Surveillance System.

Table 4 gives a breakdown of estimated submersion injuries by disposition. Children younger than 15 years of age were admitted to the hospital or treated and transferred to another hospital 43 percent of the time. For *DOA or died in the emergency department* percentages, drowning victims younger than 5 years comprised a majority of all child drownings. The deaths recorded in NEISS are also included in the fatality count in the section on reported fatalities. In contrast, for all consumer products in the CPSC’s jurisdiction, of those treated or examined in an emergency department for a product-related injury, only 4 percent of children in the younger than 15 years of age category were either admitted to the hospital or treated and transferred.

Table 4
Percent of Estimated Emergency Department-Treated Pool or Spa Submersion Injuries
Children Younger than 15 Years of Age by Disposition, 2014-2016

Disposition	Estimated Emergency Department-Treated Injury Percentages ⁷		
	Younger than 5 Years	5–14 Years	Younger than 15 Years
Examined or Treated and Released	51	53	52
Admitted to Hospital	39	34	38
Treated and Transferred	4	12	6
DOA or Died in Emergency Department	4	1	3
Held for Observation	2	0	1
Left Without Being Seen	1	-	0

Source: U. S. Consumer Product Safety Commission: National Electronic Injury Surveillance System. Appendix A details the methodology for data extraction.

⁷ Percentages may not add up to 100 due to rounding.

Table 5 shows the percentages of the estimated number of injuries for each age group by the type of location of the submersion incident. Overall, approximately half (48%) of the incidents involving injuries that led to emergency department visits occurred at a residence. Injured children younger than 5 years of age had the largest percentage (55%) in a residential location. Children 5 to 14 years of age had similar percentages in residential (31%) and public locations (33%).

Table 5
Percent of Estimated Emergency Department-Treated Pool or Spa Nonfatal Drowning Injuries
Children Younger than 15 Years of Age by Location, 2014-2016

Location	Estimated Emergency Department-Treated Injury Percentages ⁸		
	Younger than 5 Years	5–14 Years	Younger than 15 Years
Residential	55	31	48
Undisclosed Location	30	37	32
Public	16	33	20

Source: U. S. Consumer Product Safety Commission: National Electronic Injury Surveillance System. Appendix A details the methodology for data extraction.

⁸ Percentages may not add up to 100 due to rounding.

Reported Fatalities

On average, 356 fatalities associated with pool or spa submersions involving children younger than 15 years of age were reported to CPSC staff annually from 2012 through 2014. The years for reported injury and fatality statistics differ as a result of the lag in fatality reporting.

Reported fatality frequencies by year and age category are shown in Table 6. Seventy-seven percent of the victims of the reported pool- or spa-related, childhood submersion fatalities were younger than 5 years of age. As noted previously, victims in this age category also accounted for an average of 74 percent of the childhood submersion injuries related to pools or spas between 2014 and 2016. Cases in NEISS that were classified as DOA or died in the ED are also included in fatality case counts for their respective years.

For the 1,052 reported drowning incidents from 2012 through 2014, there were 1,033 fatalities (98 percent of the incidents) that involved one victim; 15 incidents that involved two victims; and four incidents that involved one victim who was included in the count, plus additional victims who were older than 14 years of age, and therefore, excluded from the counts.

The numbers of drownings related to pools or spas presented in the following section are based on all incidents reported to CPSC staff. These numbers are considered minimum counts only and cannot be used as generalized estimates for the U.S. population because they are derived from anecdotal data.

Table 6
Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas
Children Younger than 15 Years of Age, 2012-2014

Year ⁹	Reported Fatality Frequencies			
	Younger than 5 Years	5–9 Years	10–14 Years	Younger than 15 Years ¹⁰
Average	272	57	26	356
2014	252	59	34	346
2013	283	49	24	356
2012	282	63	20	365
Totals 2012-2014	817	171	78	1067

Source: CPSC databases including NEISS, IPII, DTHS, and INDP. Appendix A details the methodology for data extraction.

⁹ Reporting is not considered complete for 2013 and 2014. The number of reported fatalities may change in the future.

¹⁰ This category includes 1 case in 2014 where the victim's age is unknown, but inferred to be under 15 years of age.

Table 7 provides information on the interval between the submersion incident and the time of death for pool- or spa-related drownings. For most of the fatalities (78 percent), the date of death was either the same as the date of the incident, or 1 day later. However, 22 percent of the victims younger than 15 years of age succumbed days, weeks, and even years after the submersion, often after extensive medical treatment.

Table 7
Percentage of Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas
Children Younger than 15 Years of Age by Interval Between Injury and Death,¹¹ 2012-2014

Days Between Incident & Death	Percentage of Reported Fatalities¹²			
	Younger than 5 Years	5–9 Years	10–14 Years	Younger than 15 Years
0 days	68	71	62	68
1 day	9	12	9	10
2–7 days	16	12	23	16
8–31 days	4	4	5	4
> 31 days	2	1	1	2

Source: CPSC databases including NEISS, IPII, DTHS, and INDP. Appendix A details the methodology for data extraction.

¹¹ Note that the age at time of death is used to determine the appropriate age category. In most cases, the difference between the date of incident and date of death is not sufficient to change the age category. There were 23 fatalities where the difference was more than 31 days.

¹² Percentages may not add up to 100 due to rounding.

Reported drownings occurred predominantly in pools. A small number of drownings were associated with spas. Children younger than 5 years of age comprised almost all of the reported spa-related drownings. Table 8 records these percentages by product type.

Table 8
Percentage of Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas
Children Younger than 15 Years of Age by Product Type, 2012-2014

Product	Percentage of Reported Fatalities			
	Younger than 5 Years	5-9 Years	10-14 Years	Younger than 15 Years
Pool	95	99	100	96
Spa	5	1	-	4

Source: CPSC databases including NEISS, IPII, DTHS, and INDP. Appendix A details the methodology for data extraction.

Table 9 gives the percentages of pool or spa drownings by victim age and gender. For all age groups under 15, there were more reported male submersion victims than reported female submersion victims. This is consistent with the injury data, which show that more male children were treated in emergency departments for pool- or spa-related submersion injuries.

Table 9
Percentage of Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas Children
Younger than 15 Years of Age by Gender, 2012-2014


Gender	Percentage of Reported Fatalities ¹³			
	Younger than 5 Years	5-9 Years	10-14 Years	Younger than 15 Years
Male	66	72	81	68
Female	34	27	19	32

Source: CPSC databases including NEISS, IPII, DTHS, and INDP. Appendix A details the methodology for data extraction.

¹³ Percentages may not add up to 100 due to rounding.

Figure 3 illustrates the monthly distribution of reported pool- or spa-related childhood drownings as a function of victim age. As expected, the summer months of May, June, July, and August had the largest annual frequencies for all age groups.


Figure 3
Average Annual Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas
Children Younger than 15 Years of Age by Month of Injury and Age Category, 2012-2014



Source: CPSC databases including NEISS, IPII, DTHS, and INDP.

Figure 4 shows the annual average of reported pool or spa drownings in children younger than 15 years old as a frequency distribution of the victim's age. Children between the ages of 1 and 3 years (12 to 47 months) comprised approximately 65 percent of the reported pool or spa submersion fatalities. The graph shows a sharp decrease after age 2 (less than or equal to 35 months).

Figure 4
Average Annual Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas
Children Younger than 15 Years of Age by Age, 2012-2014



Source: CPSC databases including NEISS, IPII, DTHS, and INDP.

Table 10 records the percentages of reported pool or spa drownings by incident location. The majority of reported deaths (74 percent for pools or spas) occurred in residential settings, such as the victim’s home, a family or friend’s house, or a neighbor’s residence. The victim’s home accounts for the largest percentage (50 percent) for all location categories for victims younger than 15 years of age. For children 5 to 9 years of age and children 10 to 14 years of age, the public/community/business location accounted for the largest percentage of reported drownings.

Table 10
Percentage of Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas Children Younger than 15 Years of Age by Incident Location, 2012-2014

Location	Percentage of Reported Fatalities ¹⁴			
	Younger than 5 Years	5–9 Years	10–14 Years	Younger than 15 Years
Home	58	25	24	50
Family/ Friend	22	7	5	18
Public/ Community/ Business¹⁵	11	49	53	20
Undisclosed Location	3	16	14	6
Neighbor	7	4	4	6

Source: CPSC databases including NEISS, IPII, DTHS, and INDP. Appendix A details the methodology for data extraction.

¹⁴ Percentages may not add up to 100 due to rounding.

¹⁵ Condominium and apartment complex pools are included in this category.

Table 11 presents the percentages of reported drownings by pool/spa type. The in-ground product type accounted for the largest percentage of known pool/spa types (54 percent for victims younger than 15). This was followed by the above-ground pool category and portable pool category for cases where pool/spa type was known.

Table 11
 Percentage of Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas Children Younger than 15 Years of Age by Specific Pool/Spa Type Product Category, 2012-2014

Location	Percentage of Reported Fatalities ¹⁶			
	Younger than 5 Years	5-9 Years	10-14 Years	Younger than 15 Years
In-Ground	52	62	59	54
Undisclosed Pool/Spa Type	14	35	32	19
Above-Ground (Pools Only)	24	2	6	19
Portable ¹⁷ (Pool Only)	4	1	3	4
Inside Home (Spa Only)	1	1	-	1
Outside Home (Spa Only)	4	-	-	3

Source: CPSC databases including NEISS, IPII, DTHS, and INDP. Appendix A details the methodology for data extraction.

¹⁶ Percentages may not add up to 100 due to rounding.

¹⁷ A “portable pool” is defined as any pool that can be set up/taken down or moved to another location with relative ease.

Because the majority of reported drowning victims were younger than 5 years of age, the incident reports from 2012 through 2014 were evaluated, and common scenarios for children younger than 5 years of age for pools or spas (817 reported drownings) were classified. The highest percentage of the reports (65 percent) attributed the incident to a lapse in adult supervision (an adult losing contact or knowledge of the whereabouts of the child and, during this time period, the child managed to access the pool/spa). Ten percent of the reports indicated barrier compromise or circumvention. Another common scenario—12 percent of the reports—involved close proximity to the pool/spa, with the victim last seen in the pool/spa, or near the pool/spa, before the incident occurred. In 13 percent of the reports, there was too little information available to determine the scenario. The scenarios are categorized in Table 12. Hazard scenarios for older children are not characterized because CPSC staff receives fewer reports of drownings involving this age group.

Table 12
Percentage of Drowning Deaths Reported to CPSC Staff Associated with Pools or Spas Children Younger than 5 Years of Age by Scenario, 2012-2014

Scenario	Percentage of Reported Fatalities for Pools and Spas
Lost Contact or Knowledge of Whereabouts	65
Not Enough Information to Determine Scenario	13
Barrier Integrity or Circumvented Barrier	10
Near Pool/Spa or In Pool/Spa	12

Source: CPSC databases including NEISS, IPII, DTHS, and INDP. Appendix A details the methodology for data extraction.

Appendix A

Methodology for Pool or Spa Submersion: Estimated Nonfatal Drowning Injuries and Reported Drownings (2017)

“Drowning” is defined as suffocation and death resulting from filling of the lungs with water or other substances or fluid, so that gas exchange becomes impossible. A “non-fatal drowning” is defined as survival for any length of time after submersion in water and temporary suffocation. “Submersion” is defined as the act of placing or the condition of being under the surface of a liquid.¹⁸

Injury estimates came from NEISS data extracted on April 10, 2017, for calendar year 2016. The NEISS product codes used for the data were 3251 (Built-in pools), 3221 (Above-ground pools), 5043 (Portable pools), 1284 (Pools, not specified), 3274 (Swimming, activity) and 698 (Hot tubs and Spas). Diagnoses codes of 69 (Submersions), 65 (Anoxia), and 42 (Aspirated on) were also used, along with the age constraint of “children younger than 15 years of age,” to restrict the extracted data. Cases involving the activity of swimming were reviewed for potential inclusion in the data set. NEISS data from 2014 and 2015 were also used from last year’s report to cover the 2014 through 2016 timeframe. NEISS data is from a probability-based sample. Sampling weights are used to project the cases from NEISS hospitals to national estimates. Because incidents in NEISS are unique, there were no duplicates.

The estimated numbers of emergency department-treated injuries are rounded to the nearest hundred. Percentages in this report are rounded to the nearest integer. Because NEISS is a weighted sample, injury category percentages were based on the category weighted estimate (not rounded), divided by the total weighted estimate (not rounded).

Data were extracted on March 16, 2017, from NEISS, IPII, DTHS and INDP for pool- or spa-related submersion deaths involving children younger than 15 years of age for the years 2012 to 2014. These data were merged with data from last year’s report for 2012 and 2013, to cover the 2012 through 2014 reporting period. It should be noted that for a given year, incidents are included on an ongoing basis for IPII and DTHS. In particular, additional reports for prior reported years are generally received for the most recent years. Fatal incidents associated with product codes 3251 (Built-in pools), 3221 (Above-ground pools), 5043 (Portable pools), 1284 (Pools, not specified), 3274 (Swimming, activity), and 698 (Hot tubs and Spas) were examined for inclusion in counts. Information from these cases was extracted into an Excel spreadsheet and sorted by date and incident location. As pool submersion incidents are notable events in the community where they occur, there were often multiple news reports (IPII), a medical examiner’s report (IPII), a death certificate (DTHS), an in-depth investigation (INDP), and less frequently, a hospital emergency department report (NEISS) for a single incident. IPII is a mixture of various types of information, including newspaper clippings, consumer complaints, and reports from other government agencies, such as medical examiners/coroners. Information is submitted voluntarily to IPII, so staff cannot be sure that information on all the deaths has been received. Source documents were checked to eliminate duplicate incident reports.

¹⁸ *Dorland’s Illustrated Medical Dictionary*, 30th Edition, Saunders, 2003.